## CONTENTS

Abstract
Introduction
Interpretation of kinetic experiments in a flow system
Experimental
Apparatus
Catalyst composition and pretreatment
Synthesis tests
Kinetics of the Fischer-Tropsch synthesis on iron catalysts
Effect of space velocity, feed composition, and temperature
Pressure dependence of rate
Effect of variables of operation on selectivity
Effect of water vapor, carbon dioxide, ethane, and argon on rate
A semifundamental rate equation for the Fischer-Tropsch synthesis on iron catalysts
Rate as a function of catalyst geometry
Effect of particle size, temperature of reduction, and extent of reduction Application of a modified Wheeler equation
List of references
Appendix.—Synthesis tests on iron catalysts at pressures up to 103 atmospheres

## **ILLUSTRATIONS**

Fig	
_	Water feed system
2	Evaluation of data for 1H   100 food may 4 2400 C
3	Evaluation of data for 1H <sub>2</sub> +1CO feed gas at 240° C, reduced catalyst  Evaluation of data for 2H <sub>2</sub> +1CO feed gas at 240° C, nitrided catalyst  Evaluation of data for 2H <sub>2</sub> +1CO feed gas at 240° C, nitrided catalyst
4.	Evaluation of data for $1H_2 + 1CO$ feed gas at 240° C, nitrided catalyst
5.	
6.	Data for nitrided catalysts at 240° C according to empirical equation
7.	Plots of equation (10) for reduced catalysts
8.	Plots of first order empirical equation for nitrided catalyst D3001 at 240° C with
•	$1n_2+100 \text{ gas}_{}$
	Logarithmic plots of $(xS)_x$ against operating pressure for tests with $1H_2+1CO$ at $240^{\circ}$ C.
	Logarithmic plots of differential reaction rate $r$ as a function of partial pressure of $H_2+CO$ at $240^{\circ}$ C.
	Relative usage of hydrogen as a function of conversion for nitrided catalyst D3001 at 21.4 atmospheres and 225° C
	Methane production as a function of conversion for nitrided catalyst D3001 at 21.4 atmospheres and 225° C.
	Olefins in C <sub>2</sub> and C <sub>3</sub> fractions as a function of conversion for nitrided catalyst D3001 at 225° C and 21.4 atmospheres
	Selectivity as a function of feed gas composition, at 21.4 atmospheres and conversion of 30 percent
15.	Relative usage of hydrogen as a function of conversion and operating procesure
10.	atmospheres with 1H <sub>0</sub> +1CO during initial period of synthesis at 240° C and 21.4
	Kinetic data for synthesis on nitrided iron catalyst with 1H <sub>2</sub> +1CO gas + 10 mole-percent H <sub>2</sub> O at 21.4 atmospheres and 240° C
	Ametic data for synthesis on nitrided iron catalyst with 1H <sub>2</sub> +1CO gas + 30 mole- percent H <sub>2</sub> O at 21.4 atmospheres and 240° C
19.	Double logarithmic plots of conversion of H <sub>2</sub> +CO and CO as a function of reciprocal space velocity in H <sub>2</sub> O addition tests with 1H <sub>2</sub> +1CO at 21.4 atmospheres
	and 240 C
	Relative usage of hydrogen as a function of conversion and added water at 21.4 atmospheres and 240° C
21.	Production of $CO_2$ as a function of conversion and added water at 21.4 atmospheres and 240° $C$

<ul> <li>22. Production of methane as a function of conversion and added water vapor a 240° C and 21.4 atmospheres.</li> <li>23. Double logarithmic plots of activity against partial pressure of 1H<sub>2</sub>+1CO in feer in dilution tests.</li> <li>24. Differential reaction rates obtained by numerical differentiation.</li> <li>25. Plots of equation (29) for nitrided fused iron catalyst D3001 at 21.4 atmospheres.</li> <li>26. Variation of activity with particle size.</li> <li>27. Log-log plot of data for reduced and reduced-and-nitrided catalysts in particle size series.</li> </ul>
240° C and 21.4 atmospheres.  23. Double logarithmic plots of activity against partial pressure of 1H <sub>2</sub> +1CO in feed in dilution tests.  24. Differential reaction rates obtained by numerical differentiation.  25. Plots of equation (29) for nitrided fused iron catalyst D3001 at 21.4 atmospheres.  26. Variation of activity with particle size.  27. Log-log plot of data for reduced and reduced-and-nitrided catalysts in particle size series.
4. Differential reaction rates obtained by numerical differentiation 5. Plots of equation (29) for nitrided fused iron catalyst D3001 at 21.4 atmospheres. 6. Variation of activity with particle size 7. Log-log plot of data for reduced and reduced-and-nitrided catalysts in particle size series
Differential reaction rates obtained by numerical differentiation. Differential reaction (29) for nitrided fused iron catalyst D3001 at 21.4 atmospheres. Unitation of activity with particle size. Log-log plot of data for reduced and reduced-and-nitrided catalysts in particle size series.
<ul> <li>b. Plots of equation (29) for nitrided fused iron catalyst D3001 at 21.4 atmospheres</li> <li>c. Variation of activity with particle size</li></ul>
<ul> <li>Variation of activity with particle size</li></ul>
size series
Size series
0 0-1 4:
3. Schematic representation of a pore in a Fischer-Tropsch catalyst
9. Changes in catalyst composition in 6 weeks of synthesis expressed as change o atom ratios to iron of oxygen, carbon, and nitrogen
). Idealized representation of partly reduced particle of catalyst
. Logarithmic plots of equation (34)
2. Data for particle size series superimposed on curves of figure 31
3. Data for extent of reduction series superimposed on curves of figure 31.
Log-log plots of pressure dependence of synthesis on nitrided iron catalysts a high pressure
Distribution of hydrocarbons from synthesis with nitrided D3001 at high pres-
sures
TABLES
1. Basic data as computed from gas flows and compositions. Nitrided catalysts
with 1H <sub>2</sub> +1CO gas at 240° C
2. Values of initial reaction rate, $r_0$
iron catalysts
4. Comparison of results of iron and cobalt catalysts
o. Illiual reaction rate as a function of water added to feed
5. Constants of equation (29) for synthesis on nitrided fused from catalyst D2001 at
21.4 atmospheres 7. Variation of activity with particle size
o. Variation of activity with extent of reduction
9. Variation of activity with temperature of reduction
<ol> <li>Activation energies as a function of particle size for nitrided iron catalysts</li> <li>Values of k and D</li> </ol>
2. Values of $\gamma = D/T$
1. Fischer-1ropsen synthesis with nitrided from catalyst 1)3001 at high processing
4. I ISCHET-I TODSCH Synthesis with nitrided iron catalyst D2008 at high programme
3. Fischer-Tropsch synthesis with reduced iron catalyst D3001 at high pressures.
<ul> <li>Basic data as computed from gas flows and compositions. Nitrided catalysts with 1H<sub>2</sub>+1CO gas at 240° C.</li> <li>Values of initial reaction rate, r<sub>o</sub>.</li> <li>Activation energies from ratios of xS values at constant values of x for nitrided-iron catalysts.</li> <li>Comparison of results of iron and cobalt catalysts.</li> </ul>